Scientific Excellence • Resource Protection & Conservation • Benefits for Canadians
Excellence scientifique • Protection et conservation des ressources • Bénéfices aux Canadiens

# Oceanographic and Biological Data Relating to Short-Finned Squid (*illex illecebrousus*) *Southern Florida, January 1986*

L. Petrie

Physical and Chemical Sciences Branch
Scotia-Fundy Region
Department of Fisheries and Oceans

Bedford Institute of Oceanography
P.O. Box 1006
Dartmouth, Nova Scotia
Canada B2Y 4A2

1992

**Canadian Data Report of Hydrography and Ocean Sciences No. 111**

## Canadian Data Report Of Hydrography and Ocean Sciences

Data reports provide a medium for the documentation and dissemination of data in a form directly useable by the scientific and engineering communities. Generally, the reports contain raw and/or analyzed data but will not contain interpretations of the data. Such compilations commonly will have been prepared in support of work related to the programs and interests of the Ocean Science and Surveys (OSS) sector of the Department of Fisheries and Oceans.

Data reports are not intended for general distribution and the contents must not be referred to in other publications without prior written authorization from the issuing establishment. The correct citation appears above the abstract of each report. Data reports are abstracted in *Aquatic Sciences and Fisheries Abstracts* and indexed in the Department's annual index to scientific and technical publications.

Data reports are produced regionally but are numbered nationally. Requests for individual reports will be filled by the issuing establishment listed on the front cover and title page. Out of stock reports will be supplied for a fee by commercial agents.

Regional and headquarters establishments of Ocean Science and Surveys ceased publication of their various report series as of December 1981. A complete listing of these publications is published in the *Canadian Journal of Fisheries and Aquatic Sciences*, Volume 39: Index to Publications 1982. The current series, which begins with report number 1, was initiated in January 1982.

## Rapport statistique canadien sur l'hydrographie et les sciences océaniques

Les rapports statistiques servent de véhicule pour la compilation et la diffusion des données sous une forme directement utilisable par les scientifiques et les techniciens. En général, les rapports contiennent des données brutes ou analysées, mais ne fournissent pas d'interprétation des données. Ces compilations sont préparées le plus souvent à l'appui de travaux liés aux programmes et intérêts du service des Sciences et levés océaniques (SLO) du ministère des Pêches et des Océans.

Les rapports statistiques ne sont pas destinés à une vaste distribution et leur contenu ne doit pas être mentionné dans une publication sans une autorisation écrite préalable de l'établissement auteur. Le titre exact paraît au-dessus du résumé de chaque rapport. Les rapports statistiques sont résumés dans la revue *Résumés des sciences halieutiques et aquatiques*, et ils sont classés dans l'index annuel des publications scientifiques et techniques du Ministère.

Les rapports statistiques sont produits à l'échelon régional, mais numérotés à l'échelon national. Les demandes de rapports seront satisfaites par l'établissement auteur dont le nom figure sur la couverture et la page du titre. Les rapports épuisés sont fournis contre rétribution par des agents commerciaux.

Les établissements des Sciences et levés océaniques dans les régions et à l'administration centrale ont cessé de publier leurs diverses séries de rapports en décembre 1981. Une liste complète de ces publications figure dans le volume 39, Index des publications 1982, du *Journal canadien des sciences halieutiques et aquatiques*. La série actuelle a commencé avec la publication du rapport numéro 1 en janvier 1982.

Canadian Data Report of
Hydrography and Ocean Sciences No. 111

1992

# OCEANOGRAPHIC AND BIOLOGICAL DATA RELATING TO SHORT-FINNED SQUID (*ILLEX ILLECEBROSUS*) SOUTHERN FLORIDA, JANUARY 1986

by

L. Petrie

Physical and Chemical Sciences Branch
Scotia-Fundy Region
Department of Fisheries and Oceans

Bedford Institute of Oceanography
P.O.Box 1006
Dartmouth, Nova Scotia
Canada B2Y 4A2

©Minister of Supply and Services Canada 1992
Cat. No. FS 97-16/111E ISSN 0711-6721

Correct citation for this publication:

Petrie, L. 1992. Oceanographic and biological data relating to short-finned squid (*Illex Illecebrosus)* southern Florida, January 1986. Can. Data Rep. Hydrogr. Ocean Sci. 111: vii + 47 pp.

92-05914/



TABLE OF CONTENTS


| | |
|---|---|
| Abstract/Résumé | iv |
| List of Tables | v |
| List of Figures | vi |
| Introduction | 1 |
| Methods and Sampling | 2 |
| Data Presentation | 4 |
| Acknowledgements | 5 |
| References | 5 |
| Tables | 6 |
| Figures | 22 |

## ABSTRACT

Petrie, L. 1992. Oceanographic and biological data relating to short-finned squid (*Illex illecebrosus*), southern Florida, January 1986. Can. Data Rep. Hydrog. Ocean Sci. 111: vii + 47 pp.

Measurements of abundance, distribution and biological characteristics of larval and juvenile *Illex illecebrosus* taken in January 1986 are presented together with temperature (T), salinity (S), sigma-theta ($\sigma_\theta$), nutrients and chlorophyll-a in the vicinity of the western edge of the Gulf Stream off the Atlantic coast of Florida between 25°25'N and 29°00'N latitude.



## RÉSUMÉ

Petrie, L. 1992. Oceanographic and biological data relating to short-finned squid (*Illex illecebrosus*), southern Florida, January 1986. Can. Data Rep. Hydrog. Ocean Sci. xxx: vi + 47 pp.

On présente des mesures de l'abondance, de la distribution et des caractéristiques biologiques de *Illex illecebrosus* à l'état larvaire et à l'état juvénile, prises en janvier 1986. On présente également des données sur la temperature (T), sur la salinité (S), sur sigma-theta ($\sigma_\theta$), sur les matières nutritives et sur la chlorophylle-a aux alentours de limite ouest du Gulf Stream au large de la côte atlantique de Floride, entre 25°25' et 29°00' de latitude Nord.

# LIST OF TABLES


| | | Page |
|---|---|---|
| Table I | Station infirmation and sampling summary | 6 |
| Table II | Catches of larvae/early juveniles (by type) | 11 |
| Table III | Type A larvae/early juveniles (catch-lengths) | 14 |
| Table IV | Type B larvae/early juveniles (catch-lengths) | 17 |
| Table V | Type C larvae/early juveniles (catch-lengths) | 19 |
| Table VI | Adult squid captured (sex,catch-length,maturity stage) | 21 |

# LIST OF FIGURES


| | | Page |
|---|---|---|
| Figure 1 | Map depicting area of operation | 22 |
| Figure 2 | Transect I cross-section diagrams of temperature, salinity, sigma-θ (Stn. 1-12, CTDs 1-12) | 23 |
| Figure 3 | Transect I TS diagram (Stn. 1-12, CTDs 1-12) | 24 |
| Figure 4 | Line **N** cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram (Stn. 38-47, CTDs 13-22) | 25 |
| Figure 5 | Line **N** cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram (Stn. 47-51, CTDs 22-26) | 26 |
| Figure 6 | Line **N** cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram (Stn. 51-57, CTDs 26-32) | 27 |
| Figure 7 | Line **N** cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram (Stn. 57-62, CTDs 32-37) | 28 |
| Figure 8 | Line **N** cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram (Stn. 62-66, CTDs 37-41) | 29 |
| Figure 9 | Line **N** cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram (Stn. 66-73, CTDs 41-48) | 30 |
| Figure 10 | Line **N** cross-section diagrams of phosphate, nitrate, silicate and chlorophyll-a (Stn. 66-73, CTDs 41-48) | 31 |
| Figure 11 | Line **C** cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram (Stn. 74-81, CTDs 49-56) | 32 |
| Figure 12 | Line **C** cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram (Stn. 81-88, CTDs 56-63) | 33 |
| Figure 13 | Line **C** cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram (Stn. 88-94, CTDs 63-69) | 34 |
| Figure 14 | Line **C** cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram (Stn. 94-100, CTDs 69-75) | 35 |
| Figure 15 | Line **C** cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram (Stn. 100-106, CTDs 75-81) | 36 |
| Figure 16 | Line **C** cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram (Stn. 106-112, CTDs 81-87) | 37 |
| Figure 17 | Line **C** cross-section diagrams of phosphate, nitrate, silicate and chlorophyll-a (Stn. 106-112, CTDs 81-87) | 38 |

LIST OF FIGURES continued


| | | Page |
|---|---|---|
| Figure 18 | Line S cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram (Stn. 113-117, CTDs 88-92) | 39 |
| Figure 19 | Line S cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram (Stn. 117-121, CTDs 92-96) | 40 |
| Figure 20 | Line S cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram (Stn. 121-125, CTDs 96-100) | 41 |
| Figure 21 | Line S cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram (Stn. 125-129, CTDs 100-104) | 42 |
| Figure 22 | Line S cross-section diagrams of phosphate, nitrate, silicate and chlorophyll-a (Stn. 125-129, CTDs 100-104) | 43 |
| Figure 23 | Maps showing surface location of Slope/Gulf Stream front (solid line) on 5 occasions in the 14-28 January 1986 period. Lines N, C and S are also indicated. The location of the front was extracted from NOAA/NOS satellite-derived oceanographic analysis maps. | 44 |
| Figure 24 | Diagrams showing size distribution for Type A larvae/ early juveniles from lines N, C and S. | 45 |
| Figure 25 | Diagrams showing size distribution for Type B larvae/ early juveniles from lines N, C and S. | 46 |
| Figure 26 | Diagrams showing size distribution for Type C larvae/ early juveniles from lines N, C and S. | 47 |

## INTRODUCTION

This report presents biological and physical oceanographic data collected during R/V Alfred H. Needler cruise number 56, off south Florida, during 8-28 January, 1986. This was one in a series of cruises to determine the abundance, distribution, and types of larval and juvenile short-finned squid (*Illex Illecebrosus*) in the frontal zone between Gulf Stream Water and Slope Water from Chesapeake Bay to Florida. An additional goal was to determine the role oceanographic processes, including the Gulf Stream, play in the squid distribution and transport. Two other cruises in this series were C.S.S. Hudson Cruise 84-049, sampling from Jacksonville, Florida to Cape Hatteras from 11-14 December, 1984, and R/V Alfred H. Needler Cruise 85-001, sampling from southern Florida to Cape Hatteras from 7-22 January, 1985 (Trites and Rowell, 1985).

### R/V Alfred H. Needler Cruise Number 56

After departing Halifax, Nova Scotia, the ship steamed to approximately 29°00'N, 76°30'W and commenced Transect I, a line of 12 stations running southwestward to 27°14'N, 79°58'W. The sampling at each station consisted of an oblique bongo net tow, a CTD cast and a mid-water trawl. XBTs were released between stations, beginning after station 6 (Table I).

At the shoreward end of Transect I, a series of 15 bottom trawls were conducted in the area roughly bounded by 27°15'N to 27°50'N and 79°49'W to 80°10'W (Fig. 1). Hydrographic sampling (bottle casts) or XBTs were done at all but two of these trawl stations (Table I).

Following the bottom trawling phase, an effort was made to locate a patch of water bearing a high concentration of larvae and/or juveniles. Ten stations were occupied on two separate tracks and consisted of oblique bongo net tows, mid-water trawls, and XBTs (Fig.1, Table I).

The majority of sampling took place during multiple occupations of three transect lines. The first, ( **N** ), was located north of Cape

Canaveral along 29°00'N, from 79°20'W to 80°20'W. The second was line ( C ), west of Grand Bahama Island along 26°30'N, from 79°21'W to 80°00'W, and the final sampling line, ( S ), was east of Key Largo along 25°25'N, from 79°41'W to 80°04'W (Fig. 1).

Lines N and C were each occupied 6 times; line N from 17-21 January, and line C from 22-26 January . Sampling on both lines consisted of bongo net tows, CTDs, XBTs, and Tucker trawls. Additionally, during one occupation of each line, temperatures, salinities, nutrients, and chlorophylls were measured. Another occupation of each of these lines included mid-water trawling.

The sampling procedure on line S corresponded to that for lines N and C; however, line S was occupied only 4 times. Details of the measurements taken along these three lines appear in Table I.

## METHODS AND SAMPLING

### Physical Oceanography

The temperature and salinity measurements were taken using a portable, continuous profiling Guildline CTD (Model 8770) and recorded on cassette tape. The temperature resolution was ±0.02°C, ±0.04 for salinity and ±2.0 dBar for pressure. The data were processed at BIO using a Hewlett-Packard 1000 computer. Potential densities were calculated from the temperature and salinity measurements using the UNESCO 1980 formulation and are expressed in sigma-θ units. Calibration of temperature and salinity measurements was accomplished by comparison with surface bucket samples taken during each cast.

Water samples from hydrographic casts were obtained using Niskin bottles (5 l). The series of hydrographic stations during the bottom trawling phase of the cruise consisted of two bottles only, surface and near bottom, fitted with reversing thermometers and generating *in situ* measurements of both temperature and salinity. The hydrographic stations of Lines N , C , and S contained as many as ten Niskin bottles, depending on station depth. Reversing

thermometers were placed on the surface and deepest bottles. Salinity, phosphate, nitrate, silicate and chlorophyll-a samples were drawn from all bottles. The samples for nutrient analysis (phosphate, nitrate and silicate) were taken in 30 ml Nalgene vials, quickly frozen and stored at approximately -20°C. Samples were analyzed at BIO using standard methods (Strickland and Parsons, 1968) with a Technicon AutoAnalyzer II, sub-sampling the 30 ml vials to obtain triplicate results. Averaged values were used in reporting the nutrient concentrations. In cases where one of the three values differed appreciably from the other two, it was discarded. Values reported are micromolar (μM). Chlorophyll-a samples (250 ml) along with one drop of 1% magnesium carbonate ($MgCO_3$) were filtered through Millipore type HA 0.45 μm filters. The filters were then frozen pending analysis at BIO. The analysis followed standard methods (Yentsch and Menzel, 1963; Holm-Hansen *et al.*, 1965) using a Turner Model 10 Fluorometer. Values are reported in micrograms/litre (μg/l).

Biology

Biological sampling consisted of various combinations of bongo tows, Tucker trawls, mid-water trawls and bottom trawls (Table I). Standard MARMAP bongo nets of 505 μm mesh were towed obliquely with a ship's speed of 2.0-2.5 knots to maintain a 45° wire angle. Wire was deployed at 50 m/min and retrieved at 20 m/min. For all stations of sufficient depth, the tow was taken to 150 m. General Oceanics flow meters were centered in both bongo net openings.

The Tucker trawling operations were the same as those for the bongo tows. The mesh on the Tucker trawl was also 505 μm, and the net opening was 2.5 m x 2 m. It is important to note that the effective opening of the net would vary with depth and towing speed. Flow meters were not used with the Tucker trawl.

The mid-water trawls (Diamond IX) were also towed obliquely. Where depth permitted, the tows were to a maximum of 150 m. Trawl time was approximately 30 minutes.

Bottom trawling operations off Fort Pierce (Fig. 1) consisted

of 30 minute tows with a Western IIA trawl.

Processing of Biological Samples

All plankton were preserved in 5% buffered formalin for at least two days. As time permitted, the formalin was exchanged for, or the sample transferred to, 70% ethanol.

Catches from the mid-water trawls were sorted and all juvenile *Illex* and other cephalopods removed. Where time permitted, dorsal mantle lengths of up to 50 *Illex* were measured to the nearest millimeter. A random sub-sample of up to 2 litres of each mid-water catch was fixed in 10% formalin.

Juvenile *Illex* were initially fixed in 10% formalin for 3 days, then transferred to 70% ethanol. Occasionally, samples of a maximum of 50 individuals were preserved directly in 70% ethanol for aging studies.

Adult *Illex* were examined in detail and frozen. Representative samples were preserved in 70% ethanol. Gonads from mature specimens were preserved in Bouin's solution; the specimen from which they were taken was preserved in ethanol.

## DATA PRESENTATION

Detailed information on tabular and graphic presentations is available in the List of Tables and List of Figures. Some further information is presented below.

Table I presents station and sampling information for the cruise. The order of sampling at stations was normally bongo tow, hydrographic measurements and then trawling. Dates are January, 1986 and times are UCT. The start and stop times refer to the start of the first sampling procedure and the end of the last. All latitudes are north, all longitudes are west. The positions shown refer to the start of the first sampling procedure. The reader should be aware that a difference of several nautical miles between the start and end positions of sampling were not uncommon.

Table II shows catches of squid (labelled A, B, C and

**UNIDENTIFIED** types of larvae/early juveniles). Type **C** is *Illex Illecebrosus.* Also presented are some larvae/early juveniles identified as **A-B** (definitely not **C**) and some as **B-C** (definitely not **A**). Tables III through V present lengths of those samples which were identified by type. Lengths that appear in parentheses represent early juveniles.

In Figures 3-8, 10-15 and 17-20; which are cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram, the numbers on the diagrams refer to sequential CTD numbers, as presented in Table I.

Figures 23-25 show the size distribution of clearly identified larvae and early juveniles. In viewing these diagrams it is important for the reader to remember that lines **N** and **C** were sampled 6 times and line **S** was sampled only 4 times.

**ACKNOWLEDGEMENTS**

I wish to thank T. Rowell, chief scientist of the cruise. A special thank you to R.W. Trites, K.F. Drinkwater and B.D. Petrie.

**REFERENCES**

Holm-Hansen, O., C.J. Lorenzen, R.W. Holmes and J.D.H. Strickland. 1965. Fluorometric determination of chlorophyll. J. Cons. Int. Explor. Mer. 30: 3-15.

Trites, R.W., and T.W. Rowell. MS 1985. Larval and juvenile distribution of the short-finned squid (*Illex illecebrosus)* in the Cape Hatteras-Florida Straits area in the December-January period, 1984-1985. NAFO SCR Doc., 85/69, Serial No. N1021, 36 p.

Strickland, J.D.H., and T.R. Parsons. 1968. A practical handbook of seawater analysis. Bull. Fish. Res. Bd. Canada No. 167: 311 p.

Yentsch, C.S., and D.W. Menzel. 1963. A method for the determination of phytoplankton chlorophyll and phaeophytin by fluorescence. Deep Sea Res. 10: 221-231.

## TABLE I

### STATION INFORMATION AND SAMPLING SUMMARY

| STN# | DATE | START | STOP | LATITUDE | LONGITUDE | CTD | XBT | HYD | BON | TUC | MWT | BTM |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | TRANSECT I | | | | | | | | | | |
| 1 | 13 | 08:24 | 10:32 | 29°00.07′ | 76°30.05′ | 1 | | | X | | X | |
| 2 | 13 | 13:50 | 15:27 | 28°44.01′ | 76°59.88′ | 2 | | | X | | X | |
| 3 | 13 | 19:11 | 21:06 | 28°28.80′ | 77°29.65′ | 3 | | | X | | X | |
| 4 | 14 | 00:57 | 02:37 | 28°13.25′ | 77°59.60′ | 4 | | | X | | X | |
| 5 | 14 | 05:58 | 07:46 | 27°51.89′ | 78°30.00′ | 5 | | | X | | X | |
| 6 | 14 | 10:35 | 12:12 | 27°41.61′ | 79°00.21′ | 6 | | | X | | X | |
| | 14 | 13:20 | | 27°39.07′ | 79°02.20′ | | 1 | | | | | |
| 7 | 14 | 13:58 | 15:27 | 27°36.03′ | 79°08.42′ | 7 | | | X | | X | |
| | 14 | 15:55 | | 27°34.78′ | 79°14.38′ | | 2 | | | | | |
| 8 | 14 | 16:35 | 18:11 | 27°30.66′ | 79°18.66′ | 8 | | | X | | X | |
| | 14 | 18:42 | | 27°28.14′ | 79°25.16′ | | 3 | | | | | |
| 9 | 14 | 19:17 | 20:53 | 27°25.94′ | 79°28.95′ | 9 | | | X | | X | |
| | 14 | 21:30 | | 27°22.74′ | 79°34.89′ | | 4 | | | | | |
| 10 | 14 | 22:11 | 23:45 | 27°20.85′ | 79°38.48′ | 10 | | | X | | X | |
| | 15 | 00:05 | | 27°17.74′ | 79°48.92′ | | 5 | | | | | |
| 11 | 15 | 01:24 | 01:39 | 27°15.96′ | 79°47.53′ | 11 | | | X | | X | |
| | 15 | 04:06 | | 27°13.73′ | 79°53.20′ | | 6 | | | | | |
| 12 | 15 | 04:41 | 06:00 | 27°13.74′ | 79°57.76′ | 12 | | | X | | X | |
| | | BOTTOM TRAWLING | | | | | | | | | | |
| 13 | 15 | 08:50 | 09:20 | 27°14.90′ | 80°02.48′ | | | X | | | | X |
| 14 | 15 | 10:04 | 10:34 | 27°13.41′ | 79°49.35′ | | | X | | | | X |
| 15 | 15 | 11:44 | 12:14 | 27°20.81′ | 79°48.56′ | | | X | | | | X |
| 16 | 15 | 14:25 | 14:55 | 27°20.65′ | 80°03.32′ | | | X | | | | X |
| 17 | 15 | 15:40 | 16:10 | 27°27.15′ | 80°03.11′ | | | X | | | | X |
| 18 | 15 | 18:11 | 18:41 | 27°27.94′ | 79°48.76′ | | | X | | | | X |
| 19 | 15 | 19:47 | 20:18 | 27°32.97′ | 79°49.67′ | | | X | | | | X |
| 20 | 15 | 22:21 | 22:51 | 27°33.94′ | 79°59.74′ | | 7 | X | | | | X |
| 21 | 15 | 23:50 | 00:20 | 27°34.71′ | 80°08.27′ | | | | | | | X |
| 22 | 16 | 00:54 | 01:24 | 27°39.82′ | 80°09.78′ | | 8 | | | | | X |

**TABLE I** continued

| STN# | DATE | START | STOP | LATITUDE | LONGITUDE | CTD | XBT | HYD | BON | TUC | MWT | BTM |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 23 | 16 | 02:16 | 02:46 | 27°40.64′ | 79°59.82′ | | 9 | | | | | X |
| 24 | 16 | 04:20 | 04:50 | 27°40.80′ | 79°50.07′ | | 10 | | | | | X |
| 25 | 16 | 05:47 | 06:17 | 27°47.85′ | 79°49.04′ | | 11 | | | | | X |
| 26 | 16 | 07:48 | 08:18 | 27°47.99′ | 80°00.09′ | | 12 | | | | | X |
| 27 | 16 | 09:25 | 09:55 | 27°48.14′ | 80°09.06′ | | | | | | | X |
| | | LARVAL/JUVENILE SEARCH TRACK | | | | | | | | | | |
| 28 | 16 | 12:15 | 13:26 | 27°57.11′ | 79°59.18′ | | 13 | | X | | X | |
| 29 | 16 | 14:19 | 15:24 | 27°56.93′ | 79°54.16′ | | 14 | | X | | X | |
| 30 | 16 | 16:25 | 17:44 | 27°58.65′ | 79°49.32′ | | 15 | | X | | X | |
| 31 | 16 | 19:55 | 21:09 | 28°00.06′ | 79°44.09′ | | 16 | | X | | X | |
| 32 | 16 | 23:49 | 00:59 | 28°23.46′ | 79°30.20′ | | 17 | | X | | X | |
| 33 | 17 | 02:15 | 03:21 | 28°25.25′ | 79°34.66′ | | 18 | | X | | X | |
| 34 | 17 | 04:03 | 05:21 | 28°27.45′ | 79°40.52′ | | 19 | | X | | X | |
| 35 | 17 | 06:07 | 07:27 | 28°29.21′ | 79°45.77′ | | 20 | | X | | X | |
| 36 | 17 | 08:08 | 09:30 | 28°31.03′ | 79°50.85′ | | 21 | | X | | X | |
| 37 | 17 | 11:11 | 12:42 | 28°33.94′ | 79°56.16′ | | 22 | | X | | X | |
| | 17 | 13:21 | | 28°34.77′ | 80°01.21′ | | 23 | | | | | |
| | | LINE N | | | | | | | | | | |
| 38 | 17 | 16:15 | 18:07 | 28°59.66′ | 80°19.58′ | 13 | | | X | X | | |
| 39 | 17 | 18:48 | 19:47 | 29°00.00′ | 80°13.88′ | 14 | | | X | X | | |
| 40 | 17 | 20:24 | 21:11 | 28°59.92′ | 80°08.45′ | 15 | | | X | X | | |
| 41 | 17 | 22:02 | 23:07 | 29°00.51′ | 80°02.94′ | 16 | | | X | X | | |
| 42 | 17 | 23:49 | 01:36 | 29°00.92′ | 79°57.13′ | 17 | | | X | X | | |
| 43 | 18 | 02:35 | 04:02 | 29°00.20′ | 79°50.76′ | 18 | | | X | X | | |
| 44 | 18 | 04:42 | 06:27 | 29°00.00′ | 79°45.22′ | 19 | | | X | X | | |
| 45 | 18 | 07:21 | 08:41 | 28°59.84′ | 79°39.36′ | 20 | | | X | X | | |
| 46 | 18 | 09:27 | 11:25 | 28°59.81′ | 79°33.59′ | 21 | | | X | X | | |
| 47 | 18 | 12:04 | 13:20 | 29°00.30′ | 79°28.33′ | 22 | | | X | X | | |
| | 18 | 13:50 | | 29°00.14′ | 79°22.95′ | | 24 | | | | | |
| 48 | 18 | 15:47 | 17:48 | 29°00.30′ | 79°45.37′ | 23 | | | X | X | | |
| | 18 | 18:15 | | 28°59.87′ | 79°47.42′ | | 25 | | | | | |
| 49 | 18 | 18:38 | 20:19 | 29°00.01′ | 79°51.15′ | 24 | | | X | X | | |
| | 18 | 20:45 | | 29°00.39′ | 79°53.70′ | | 26 | | | | | |

**TABLE I** continued

| STN# | DATE | START | STOP | LATITUDE | LONGITUDE | CTD | XBT | HYD | BON | TUC | MWT | BTM |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 50 | 18 | 21:03 | 23:07 | 29°00.04′ | 79°56.94′ | 25 | | | X | X | | |
| | 18 | 23:20 | | 29°00.64′ | 80°00.96′ | | 27 | | | | | |
| 51[1] | 18 | 23:31 | 00:57 | 29°00.24′ | 80°02.48′ | 26 | | | X | X | | |
| 52 | 19 | 01:27 | 03:45 | 29°00.13′ | 79°57.34′ | 27 | | | X | X | | |
| 53 | 19 | 04:30 | 06:16 | 29°00.04′ | 79°50.97′ | 28 | | | X | X | | |
| 54 | 19 | 06:54 | 08:43 | 28°59.92′ | 79°45.23′ | 29 | 28 | | X | X | | |
| 55[2] | 19 | 09:15 | 10:37 | 28°59.94′ | 79°51.02′ | 30 | | | X | | | |
| 56 | 19 | 12:12 | 14:10 | 29°00.35′ | 79°39.73′ | 31 | | | X | X | | |
| 57 | 19 | 14:54 | 16:26 | 29°00.22′ | 79°33.23′ | 32 | 29 | | X | X | | |
| 58 | 19 | 17:12 | 18:58 | 29°00.04′ | 79°39.55′ | 33 | | | X | X | | |
| 59 | 19 | 19:36 | 21:14 | 29°00.08′ | 79°45.40′ | 34 | | | X | X | | |
| 60 | 19 | 22:08 | 23:37 | 29°00.33′ | 79°50.92′ | 35 | | | X | X | | |
| 61 | 20 | 00:09 | 02:10 | 29°00.32′ | 79°56.83′ | 36 | 30 | | X | X | | |
| 62 | 20 | 03:30 | 06:27 | 29°00.11′ | 80°01.97′ | 37 | | | X | X | X | |
| 63 | 20 | 07:10 | 09:06 | 29°00.23′ | 79°57.14′ | 38 | | | X | | X | |
| 64 | 20 | 10:07 | 12:08 | 29°00.73′ | 79°51.34′ | 39 | | | X | | X | |
| 65 | 20 | 12:57 | 14:47 | 29°00.17′ | 79°45.38′ | 40 | | | X | | X | |
| 66 | 20 | 15:52 | 20:03 | 29°00.37′ | 79°40.32′ | 41 | | X | X | X | X | |
| 67 | 20 | 20:41 | 00:12 | 29°00.40′ | 79°45.84′ | 42 | | X | X | X | X | |
| 68 | 21 | 01:02 | 05:21 | 29°00.31′ | 79°51.08′ | 43 | | X | X | X | X | |
| 69 | 21 | 05:59 | 09:26 | 29°01.37′ | 79°57.62′ | 44 | | X | X | X | X | |
| 70 | 21 | 10:25 | 16:41 | 29°00.17′ | 80°02.41′ | 45 | | X | X | X | X | |
| 71 | 21 | 16:24 | 17:18 | 29°01.10′ | 80°08.67′ | 46 | | X | X | X | | |
| 72 | 21 | 18:14 | 19:06 | 29°00.40′ | 80°14.07′ | 47 | | X | X | X | | |
| 73[3] | 21 | 19:52 | 20:43 | 29°00.40′ | 80°19.93′ | 48 | | X | X | X | | |
| | | LINE C | | | | | | | | | | |
| 74 | 22 | 18:11 | 19:07 | 26°30.00′ | 79°59.97′ | 49 | | | X | X | | |
| 75 | 22 | 19:40 | 21:10 | 26°30.03′ | 79°54.33′ | 50 | | | X | X | | |
| 76 | 22 | 21:58 | 23:55 | 26°30.92′ | 79°48.25′ | 51 | | | X | X | | |
| 77 | 23 | 00:19 | 01:54 | 26°30.32′ | 79°43.59′ | 52 | | | X | X | | |
| 78 | 23 | 02:33 | 04:04 | 26°30.54′ | 79°38.52′ | 53 | | | X | X | | |

1 TUCKER TRAWL, BONGO TOW ORDER REVERSED THIS SET

2 BRIDGE HEADED WEST IN ERROR ON THIS W →→ E TRANSECT

3 TUCKER TRAWL, BONGO TOW ORDER REVERSED THIS SET

**TABLE I** continued

| STN# | DATE | START | STOP | LATITUDE | LONGITUDE | CTD | XBT | HYD | BON | TUC | MWT | BTM |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 79 | 23 | 04:43 | 06:36 | 26°30.00′ | 79°32.60′ | 54 | | | X | X | | |
| 80 | 23 | 06:57 | 08:32 | 26°30.02′ | 79°27.28′ | 55 | | | X | X | | |
| 81 | 23 | 09:07 | 10:36 | 26°30.03′ | 79°21.44′ | 56 | | | X | X | | |
| 82 | 23 | 11:08 | 12:46 | 26°30.44′ | 79°27.76′ | 57 | | | X | X | | |
| 83 | 23 | 13:23 | 14:46 | 26°30.37′ | 79°33.27′ | 58 | | | X | X | | |
| 84 | 23 | 15:12 | 16:29 | 26°30.04′ | 79°37.77′ | 59 | | | X | X | | |
| 85 | 23 | 17:16 | 19:04 | 26°30.56′ | 79°43.49′ | 60 | | | X | X | | |
| 86 | 23 | 19:46 | 21:38 | 26°30.52′ | 79°48.03′ | 61 | | | X | X | | |
| 87 | 23 | 22:18 | 23:45 | 26°30.28′ | 79°54.23′ | 62 | | | X | X | | |
| 88 | 24 | 00:16 | 01:32 | 26°30.13′ | 79°59.65′ | 63 | | | X | X | | |
| 89 | 24 | 01:58 | 03:42 | 26°30.09′ | 79°54.78′ | 64 | | | X | X | | |
| 90 | 24 | 04:25 | 05:47 | 26°30.05′ | 79°47.90′ | 65 | | | X | X | | |
| 91 | 24 | 06:19 | 08:51 | 26°30.01′ | 79°43.35′ | 66 | | | X | X | | |
| 92 | 24 | 08:27 | 10:11 | 26°29.89′ | 79°38.07′ | 67 | | | X | X | | |
| 93 | 24 | 10:40 | 12:21 | 26°30.32′ | 79°32.88′ | 68 | | | X | X | | |
| 94 | 24 | 12:50 | 14:31 | 26°30.02′ | 79°27.68′ | 69 | | | X | X | | |
| 95 | 24 | 15:11 | 17:07 | 26°32.50′ | 79°35.04′ | 70 | 31 | | X | X | | |
| 96 | 24 | 17:39 | 19:08 | 26°29.91′ | 79°38.40′ | 71 | | | X | X | | |
| 97 | 24 | 19:39 | 21:16 | 26°29.74′ | 79°43.57′ | 72 | | | X | X | | |
| 98 | 24 | 22:02 | 23:49 | 26°29.77′ | 79°47.42′ | 73 | | | X | X | | |
| 99 | 25 | 00:16 | 01:48 | 26°30.18′ | 79°54.32′ | 74 | | | X | X | | |
| 100 | 25 | 02:15 | 03:25 | 26°31.37′ | 80°00.25′ | 75 | | | X | X | | |
| 101 | 25 | 03:59 | 05:19 | 26°30.23′ | 79°54.64′ | 76 | | | X | X | | |
| 102 | 25 | 05:51 | 07:36 | 26°30.30′ | 79°48.20′ | 77 | | | X | X | | |
| 103 | 25 | 08:03 | 09:51 | 26°30.28′ | 79°43.07′ | 78 | | | X | X | | |
| 104 | 25 | 10:48 | 13:45 | 26°30.26′ | 79°38.33′ | 79 | 32 | | X | X | | |
| 105 | 25 | 14:14 | 15:33 | 26°30.15′ | 79°33.12′ | 80 | | | X | X | | |
| 106 | 25 | 16:09 | 19:42 | 26°30.09′ | 79°27.47′ | 81 | | X | X | X | X | |
| 107 | 25 | 20:16 | 23:35 | 26°29.93′ | 79°32.57′ | 82 | | X | X | X | X | |
| 108 | 26 | 00:06 | 03:17 | 26°30.18′ | 79°38.04′ | 83 | | X | X | X | X | |
| 109 | 26 | 03:44 | 07:19 | 26°30.26′ | 79°42.97′ | 84 | | X | X | X | X | |
| 110 | 26 | 07:49 | 11:16 | 26°29.70′ | 79°47.82′ | 85 | | X | X | X | X | |
| 111 | 26 | 12:01 | 13:08 | 26°30.01′ | 79°54.42′ | 86 | | X | X | | | |

**TABLE I** continued

| STN# | DATE | START | STOP | LATITUDE | LONGITUDE | CTD | XBT | HYD | BON | TUC | MWT | BTM |
|---|---|---|---|---|---|---|---|---|---|---|---|---|
| 112 | 26 | 14:09 | 17:02 | 26°29.95′ | 79°59.86′ | 87 | | X | X | X | X | |
| | | LINE S | | | | | | | | | | |
| 113 | 26 | 23:13 | 00:14 | 25°25.27′ | 80°04.11′ | 88 | | | X | X | | |
| 114 | 27 | 01:07 | 03:05 | 25°24.64′ | 79°58.19′ | 89 | | | X | X | | |
| 115 | 27 | 04:03 | 05:46 | 25°24.99′ | 79°53.06′ | 90 | | | X | X | | |
| 116 | 27 | 06:34 | 08:21 | 25°25.30′ | 79°47.21′ | 91 | | | X | X | | |
| 117 | 27 | 10:03 | 11:50 | 25°25.23′ | 79°41.57′ | 92 | | | X | X | | |
| 118 | 27 | 12:17 | 14:06 | 25°25.62′ | 79°47.40′ | 93 | | | X | X | | |
| 119 | 27 | 14:36 | 16:20 | 25°25.23′ | 79°53.07′ | 94 | | | X | X | | |
| 120 | 27 | 16:58 | 18:40 | 25°25.18′ | 79°58.64′ | 95 | 33 | | X | X | | |
| 121 | 27 | 19:21 | 20:45 | 25°25.31′ | 80°04.03′ | 96 | | | X | X | | |
| 122 | 27 | 21:17 | 22:53 | 25°25.28′ | 79°58.32′ | 97 | | | X | X | | |
| 123 | 27 | 23:28 | 01:19 | 25°25.81′ | 79°52.04′ | 98 | | | X | X | | |
| 124 | 28 | 01:58 | 03:52 | 25°25.37′ | 79°46.89′ | 99 | | | X | X | | |
| 125 | 28 | 04:39 | 07:29 | 25°25.47′ | 79°41.49′ | 100 | | X | X | | X | |
| 126 | 28 | 08:32 | 11:37 | 25°25.20′ | 79°47.54′ | 101 | | X | X | | X | |
| 127 | 28 | 12:24 | 14:49 | 25°25.40′ | 79°53.03′ | 102 | | X | X | | X | |
| 128 | 28 | 15:38 | 18:08 | 25°24.81′ | 79°58.86′ | 103 | | X | X | | X | |
| 129 | 28 | 18:39 | 20:23 | 25°25.12′ | 80°03.98′ | 104 | | X | X | | X | |

## TABLE II

### CATCHES OF LARVAE / EARLY JUVENILES

| STN# | TYPE A | | TYPE B | TYPE C | UNIDENTIFIED |
|---|---|---|---|---|---|
| 30 | 1 | | | | 1 |
| 36 | 1 | | | | |
| 38 | | | 3 | 7 | |
| 40 | | | | 1 | |
| 41 | | | | | 1 |
| 42 | 7 | | 4 | 7 | 1 |
| 44 | 1 | | 2 | 1 | |
| 45 | 1 | | 1 | 1 | |
| 48 | 1 | | | 2 | |
| 49 | 5 | | 2 | 2 | 1 |
| 50 | | | 5 | | |
| 51 | 1 | | | | |
| 52 | 2 | | | 5 | 1 |
| 53 | 6 | 1[1] | 5 | 10 | 1 |
| 54 | | | 1 | | |
| 58 | 1 | | | | |
| 59 | | | | 1 | |
| 60 | 2 | | | | |
| 61 | 4 | | 5 | 5 | 5 |
| 64 | | | | | 1 |
| 66 | 1 | | 1[2] | | |
| 67 | 1 | | 2 | | 1 |
| 68 | 3 | | 3 | 2 | 1 |
| 69 | 2 | 1 | 1 | 1 | 1 |

1 NUMBERS IN THIS INTERCOLUMNAR AREA OF TABLE II REPRESENT LARVAE THAT ARE TYPE A OR B BUT DEFINITELY NOT TYPE C .LENGTHS OF "A-B" TYPES WILL APPEAR WITH TYPE A LENGTHS IN TABLE III.

2 NUMBERS IN THIS INTERCOLUMNAR AREA OF TABLE II REPRESENT LARVAE THAT ARE TYPE B OR C BUT DEFINITELY NOT TYPE A .LENGTHS OF "B-C" TYPES WILL APPEAR WITH TYPE B LENGTHS IN TABLE IV.

**TABLE II** continued

CATCHES OF LARVAE / EARLY JUVENILES

| STN# | TYPE A | | TYPE B | | TYPE C | UNIDENTIFIED |
|---|---|---|---|---|---|---|
| 70 | 17 | | 4 | | 4 | 1 |
| 71 | 3 | | 4 | | 4 | 2 |
| 72 | | | 1 | | | |
| 74 | 2 | | 1 | | 9 | 2 |
| 75 | 2 | | 1 | | | 2 |
| 76 | 1 | | 1 | | 1 | 1 |
| 77 | 4 | | 2 | | 1 | 1 |
| 78 | 3 | | 1 | | 3 | 3 |
| 79 | | | 2 | | 5 | |
| 80 | | | 2 | | | |
| 81 | 5 | | 1 | | 4 | |
| 82 | | | 1 | | | |
| 83 | 5 | 2 | | | 1 | |
| 84 | 7 | | 5 | | 2 | 5 |
| 85 | 8 | | 5 | | 8 | 4 |
| 86 | 1 | | 2 | | 2 | |
| 87 | 1 | | | | 1 | |
| 88 | | | | 1 | 12 | 6 |
| 89 | 3 | | | | 3 | |
| 90 | | | | | | 3 |
| 91 | 1 | | 2 | | | |
| 92 | 5 | | 4 | | | 1 |
| 93 | | | | | 2 | 1 |
| 94 | 5 | | | | | |
| 95 | | | | | 1 | 1 |
| 96 | | | | | 1 | |
| 97 | 6 | | 4 | | | 1 |
| 98 | 3 | | 1 | | 5 | 2 |
| 99 | | | | | 2 | |
| 100 | | | 3 | | 8 | 3 |
| 101 | 1 | | | | 3 | 1 |

**TABLE II** continued

CATCHES OF LARVAE / EARLY JUVENILES

| STN# | TYPE A | | TYPE B | TYPE C | UNIDENTIFIED |
|---|---|---|---|---|---|
| 102 | 6 | 1 | 4 | 4 | 4 |
| 103 | 1 | | 3 | 6 | |
| 104 | 5 | | 4 | 6 | 9 |
| 105 | | | 1 | | |
| 106 | 2 | | | | |
| 107 | 1 | | | | |
| 108 | 5 | | 1 | | |
| 109 | 3 | | 4 | 2 | |
| 110 | 8 | 9 | 5 | 8 | 1 |
| 111 | | | | 2 | |
| 112 | | | 1 | 16 | 3 |
| 113 | | | 1 | 5 | |
| 114 | 4 | | 4 | | 2 |
| 115 | 6 | 3 | 4 | 8 | 7 |
| 116 | 1 | | | | |
| 117 | 3 | 4 | 3 | 5 | 2 |
| 118 | 2 | | 1 | 1 | |
| 119 | 4 | | 3 | 2 | 1 |
| 120 | 5 | 11 | | 9 | 6 |
| 121 | 4 | 2 | 1 | 17 | 1 |
| 122 | 5 | 1 | 1 | | 5 |
| 123 | 2 | | 3 | | 3 |
| 125 | | | 1 | 1 | |
| 128 | | | | 1 | |

**TABLE III**

TYPE A LARVAE / EARLY JUVENILES

| STN# | | CATCH-LENGTHS (mm) | TOTAL |
|---|---|---|---|
| 30 | | 3.6 | 1 |
| 36 | | 2.6 | 1 |
| 42 | | 2.5 2.8 3.8 2.7 2.3 3.5 3.1 | 7 |
| 44 | | 2.8 | 1 |
| 45 | | 2.5 | 1 |
| 48 | | 3.7 | 1 |
| 49 | | 3.5 1.9 2.0 2.4 3.4 | 5 |
| 51 | | 2.3 | 1 |
| 52 | | 2.1 2.0 | 2 |
| 53 | | 1.9 2.0 2.3 4.2 4.2 2.5 | 6 |
| 53 | A-B | 3.0 | 1 |
| 58 | | 1.5 | 1 |
| 60 | | 2.6 3.8 | 2 |
| 61 | | 3.2 3.5 3.3 3.1 | 4 |
| 66 | | 3.0 | 1 |
| 67 | | 2.7 | 1 |
| 68 | | 4.4 2.8 3.4 | 3 |
| 69 | | 2.1 2.4 | 2 |
| 69 | A-B | 5.0 | 1 |
| 70 | | 1.7 2.2 2.6 3.2 3.4 3.6 3.8 3.8 4.0 | |
| | | 4.0 4.2 5.6 1.9 3.0 3.0 2.8 3.8 | 17 |
| 71 | | 2.3 2.4 3.8 | 3 |
| 74 | | 1.7 1.8 | 2 |
| 75 | | 2.8 3.5 | 2 |
| 76 | | 4.8 | 1 |
| 77 | | 2.4 2.5 2.2 2.8 | 4 |
| 78 | | 2.2 2.8 2.9 | 3 |
| 81 | | 2.0 3.4 3.6 3.6 4.8 | 5 |
| 83 | | 2.2 3.0 3.3 4.8 2.7 | 5 |
| 83 | A-B | 2.4 5.1 | 2 |
| 84 | | 2.0 2.3 3.0 2.5 4.0 4.0 3.8 | 7 |

**TABLE III** continued

TYPE A LARVAE / EARLY JUVENILES

| STN# | | CATCH-LENGTHS (mm) | TOTAL |
|---|---|---|---|
| 85 | | 1.9 2.0 2.2 2.6 2.8 3.5 3.4 4.4 | 8 |
| 86 | | 1.6 | 1 |
| 87 | | 5.1 | 1 |
| 89 | | 2.0 3.3 5.0 | 3 |
| 91 | | 2.3 | 1 |
| 92 | | 1.6 3.0 3.1 3.5 2.4 | 5 |
| 94 | | 3.0 3.2 3.5 3.8 4.2 | 5 |
| 97 | | 1.8 1.8 2.0 2.0 2.6 2.8 | 6 |
| 98 | | 2.3 2.6 3.2 | 3 |
| 101 | | 2.9 | 1 |
| 102 | | 2.2 1.9 3.5 4.2 4.2 5.3 | 6 |
| 102 | A-B | 3.9 | 1 |
| 103 | | 2.1 | 1 |
| 104 | | 1.7 2.0 2.2 3.5 3.7 | 5 |
| 106 | | 2.0 1.7 | 2 |
| 107 | | 3.9 | 1 |
| 108 | | 1.6 2.0 2.6 2.0 2.2 | 5 |
| 109 | | 1.4 1.9 4.2 | 3 |
| 110 | | 2.0 2.5 2.8 2.8 3.1 3.7 4.0 4.6 | 8 |
| 110 | A-B | 2.0 2.6 3.0 3.4 3.4 3.2 3.6 3.8 5.6 | 9 |
| 114 | | 3.3 3.6 5.0 5.0 | 4 |
| 115 | | 1.7 1.7 3.3 3.9 4.6 1 head | 6 |
| 115 | A-B | 2.2 2.2 3.4 | 3 |
| 116 | | 3.9 | 1 |
| 117 | | 2.5 2.0 3.0 | 3 |
| 117 | A-B | 3.0 7.5 4.4 4.7 | 4 |
| 118 | | 2.4 2.4 | 2 |
| 119 | | 2.1 2.5 3.0 4.6 | 4 |
| 120 | | 2.1 2.2 2.8 3.0 (5.4) | 5 |
| 120 | A-B | 1.8 1.8 2.0 2.2 2.6 2.8 2.8 2.9 3.4 3.6 3.6 | 11 |
| 121 | | 3.5 4.0 3.5 3.6 | 4 |

**TABLE III** continued

TYPE A LARVAE / EARLY JUVENILES

| STN# | | CATCH-LENGTHS (mm) | TOTAL |
|---|---|---|---|
| 121 | A-B | 3.5 1.8 | 2 |
| 122 | | 2.7 3.5 3.8 4.1 1 head | 5 |
| 122 | A-B | 2.4 | 1 |
| 123 | | 2.2 2.4 | 2 |

**TABLE IV**

TYPE B LARVAE / EARLY JUVENILES

| STN# | | CATCH-LENGTHS (mm) | TOTAL |
|---|---|---|---|
| 38 | | 2.0 2.8 3.3 | 3 |
| 42 | | 2.4 2.4 3.0 4.0 | 4 |
| 44 | | 3.6 5.0 | 2 |
| 45 | | 2.9 | 1 |
| 49 | | 2.3 3.2 | 2 |
| 50 | | 2.7 3.5 3.7 4.0 5.0 | 5 |
| 53 | | 2.3 2.4 2.6 3.7 2.5 | 5 |
| 54 | | 3.8 | 1 |
| 61 | | 3.6 2.0 3.0 3.2 4.2 | 5 |
| 66 | B-C | 1.5 | 1 |
| 67 | | 1.7 2.0 | 2 |
| 68 | | 4.2 4.2 4.6 | 3 |
| 69 | | 5.2 | 1 |
| 70 | | 2.6 3.0 3.4 4.8 | 4 |
| 71 | | 4.2 4.7 4.8 5.4 | 4 |
| 72 | | 4.5 | 1 |
| 74 | | 1.8 | 1 |
| 75 | | 3.1 | 1 |
| 76 | | 3.1 | 1 |
| 77 | | 4.0 4.8 | 2 |
| 78 | | 3.6 | 1 |
| 79 | | 2.6 2.8 | 2 |
| 80 | | 4.6 2.9 | 2 |
| 81 | | 5.5 | 1 |
| 82 | | 4.4 | 1 |
| 84 | | 4.0 5.4 2.0 2.5 4.0 | 5 |
| 85 | | 2.8 3.5 4.2 4.6 5.2 | 5 |
| 86 | | 4.3 4.8 | 2 |
| 88 | B-C | 4.3 | 1 |
| 91 | | 2.5 3.3 | 2 |
| 92 | | 4.8 5.2 2.0 3.0 | 4 |

**TABLE IV** continued

TYPE B LARVAE / EARLY JUVENILES

| STN# | CATCH-LENGTHS (mm) | TOTAL |
|---|---|---|
| 97 | 2.8 2.0 2.0 3.0 | 4 |
| 98 | (5.4) | 1 |
| 100 | 1.9 2.5 2.8 | 3 |
| 102 | 2.8 3.1 4.0 (5.2) | 4 |
| 103 | 1.7 4.0 3.0 | 3 |
| 104 | 1.6 1.8 2.1 2.6 | 4 |
| 105 | 4.4 | 1 |
| 108 | 3.1 | 1 |
| 109 | 2.0 4.7 3.8 (5.0) | 4 |
| 110 | 1.7 2.2 2.8 3.2 4.4 | 5 |
| 112 | 5.0 | 1 |
| 113 | 2.5 | 1 |
| 114 | 2.6 (4.5) (5.0) (5.3) | 4 |
| 115 | 1.5 2.0 2.2 3.0 | 4 |
| 117 | 2.1 3.0 2.9 | 3 |
| 118 | 4.1 | 1 |
| 119 | 1.5 2.0 2.2 | 3 |
| 121 | 1.8 | 1 |
| 122 | 3.8 | 1 |
| 123 | 1.6 2.0 2.1 | 3 |
| 125 | 3.2 | 1 |

**TABLE V**

## TYPE C LARVAE / EARLY JUVENILES

| STN# | CATCH-LENGTHS (mm) | TOTAL |
|---|---|---|
| 38 | 2.2 2.6 2.6 3.1 3.1 3.2 3.5 | 7 |
| 40 | 2.9 | 1 |
| 42 | 2.2 2.8 3.2 3.5 5.2 3.1 2.2 | 7 |
| 44 | 2.5 | 1 |
| 45 | 4.8 | 1 |
| 48 | (5.4) (7.0) | 2 |
| 49 | 2.0 3.1 | 2 |
| 52 | 1.5 1.5 1.5 1.5 1.6 | 5 |
| 53 | 2.0 2.0 2.2 2.5 2.5 2.8 2.8 3.0 2.0 2.5 | 10 |
| 59 | 3.2 | 1 |
| 61 | 2.2 3.7 4.6 2.2 1-no mantle | 5 |
| 68 | 2.6 1.6 | 2 |
| 69 | 3.0 | 1 |
| 70 | 2.0 2.2 3.2 3.9 | 4 |
| 71 | 1.5 1.6 2.2 1-no mantle | 4 |
| 74 | 2.3 1.4 1.4 1.5 1.5 1.6 1.6 1.7 1.8 | 9 |
| 76 | 2.5 | 1 |
| 77 | 4.0 | 1 |
| 78 | 1.9 2.5 (6.0) | 3 |
| 79 | 1.8 1.9 2.0 2.8 (6.5) | 5 |
| 81 | 2.4 2.3 2.5 2.8 | 4 |
| 83 | 1.5 | 1 |
| 84 | 2.3 2.1 | 2 |
| 85 | 1.6 2.4 2.4 3.0 3.2 3.4 2.5 4.0 | 8 |
| 86 | 2.3 2.0 | 2 |
| 87 | 2.5 | 1 |
| 88 | 1.5 1.8 2.0 3.0 4.5 5.2 5.6 2.5 3.2 3.5 4.5 2.0 | 12 |
| 89 | 1.6 1.9 2.0 | 3 |
| 93 | 2.0 2.0 | 2 |
| 95 | 2.0 | 1 |

**TABLE V** continued

TYPE C LARVAE / EARLY JUVENILES

| STN# | CATCH-LENGTHS (mm) | TOTAL |
|---|---|---|
| 96 | 2.1 | 1 |
| 98 | 1.8 3.2 3.8 2.6 5.0 | 5 |
| 99 | 1.6 1.8 | 2 |
| 100 | 1.5 1.7 1.8 2.0 2.4 3.8 2-dried,unmeasurable | 8 |
| 101 | 2.0 2.4 2.2 | 3 |
| 102 | 2.6 1.5 3.5 2.6 | 4 |
| 103 | 2.0 2.2 1.9 3.1 3.4 (6.3) | 6 |
| 104 | 1.5 1.5 1.6 2.1 2.3 3.0 | 6 |
| 109 | 1.9 3.5 | 2 |
| 110 | 1.3 1.4 1.7 2.4 2.5 2.6 1.4 2.3 | 8 |
| 111 | 1.7 2.1 | 2 |
| 112 | 1.9 1.9 2.1 2.2 2.3 2.3 2.4 2.8 2.9 3.0 3.3 3.8 4.4 4.4 1.8 1.7 | 16 |
| 113 | 1.7 2.0 2.4 2.6 2.0 | 5 |
| 115 | 1.5 1.5 1.7 2.1 2.2 3.0 3.0 1-unmeasurable | 8 |
| 117 | 1.5 1.8 2.2 2.6 3.0 | 5 |
| 118 | 2.2 | 1 |
| 119 | 2.2 2.6 | 2 |
| 120 | 2.1 2.2 2.8 3.0 2.1 2.2 2.3 2.5 2.8 | 9 |
| 121 | 1.8 2.0 2.0 2.0 2.0 2.2 2.2 2.4 2.4 2.5 2.7 2.8 2.8 2.9 3.0 2.8 1-head | 17 |
| 125 | 3.4 | 1 |
| 128 | 2.0 | 1 |

**TABLE VI**

ADULT SQUID CAPTURED

| STN# | SEX | CATCH-LENGTHS (mm) | MATURITY STAGE | COMMENTS |
|---|---|---|---|---|
| 7 | | | | Two illex captured, total weight 0.014 kg. |
| 15 | M | 160 | 3 | Mature |
| | F | 165 | 3 | Maturing |
| | F | 175 | 3 | Maturing |
| 18 | M | 113 | 1 | Immature |
| | M | 145 | 4 | Spent |
| | M | 159 | 4 | Spent |
| | F | 165 | 5 | Fully mature |
| 19 | M | 125 | 2 | Maturing |
| | M | 130 | 2 | Maturing |
| | F | 124 | 3 | Maturing |
| | F | 130 | 3 | Maturing |
| 24 | M | 115 | 1 | Immature |
| | M | 120 | 2 | Maturing |
| 25 | F | 165 | 5 | Fully mature |




Fig. 1. Map depicting area of operation.




Fig. 2. Transect I , Stn. 1-12, CTDs 1-12. Cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram.




Fig. 3. Transect I , Stn. 1-12, CTDs 1-12. TS diagram.




Fig. 4. Line M , Stn. 38-47, CTDs 13-22. Cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram.




Fig. 5. Line N , Stn. 47-51, CTDs 22-26. Cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram.




Fig. 6. Line N , Stn. 51-57, CTDs 26-32. Cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram.




Fig. 7. Line N , Stn. 57-62, CTDs 32-37. Cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram.




Fig. 8. Line N , Stn. 62-66, CTDs 37-41. Cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram.




Fig. 9. Line **N** , Stn. 66-73, CTDs 41-48. Cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram.




Fig. 10. Line N , Stn. 66-73, CTDs 41-48. Cross-section diagrams of phosphate, nitrate, silicate and chlorophyll-a.




Fig. 11. Line C , Stn. 74-81, CTDs 49-56. Cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram.




Fig. 12. Line C , Stn. 81-88, CTDs 56-63. Cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram.




Fig. 13. Line C , Stn. 88-94, CTDs 63-69. Cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram.




Fig. 14. Line C , Stn. 94-100, CTDs 69-75. Cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram.




Fig. 15. Line C , Stn. 100-106, CTDs 75-81. Cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram.




Fig. 16. Line C , Stn. 106-112, CTDs 81-87. Cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram.




Fig. 17. Line C , Stn. 106-112, CTDs 81-87. Cross-section diagrams of phosphate, nitrate, silicate and chlorophyll-a.




Fig. 18. Line S , Stn. 113-117, CTDs 88-92. Cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram.




Fig. 19. Line S , Stn. 117-121, CTDs 92-96. Cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram.




Fig. 20. Line S , Stn. 121-125, CTDs 96-100. Cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram.




Fig. 21. Line S , Stn. 125-129, CTDs 100-104. Cross-section diagrams of temperature, salinity, sigma-θ and a TS diagram.




Fig. 22. Line S , Stn. 125-129, CTDs 100-104. Cross-section diagrams of phosphate, nitrate, silicate and chlorophyll-a.




Fig. 23. Maps showing surface location of Slope/Gulf Stream front (solid line) on 5 occasions in the 14-28 January 1986 period. Lines N, C and S are also indicated. The location of the front was extracted from NOAA/NOS satellite-derived oceanographic analysis maps.




Fig. 24. Diagrams showing size distribution for Type A larvae/early juveniles from lines N, C and S.




Fig. 25. Diagrams showing size distribution for Type B larvae/early juveniles from lines N, C and S.




Fig. 26. Diagrams showing size distribution for Type C larvae/early juveniles from lines N, C and S.